

Howard Johnson's Viper[®] Granular Weed Control

ACTIVE INGREDIENTS:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid* 1.346%
2,4-Dichlorophenoxyacetic Acid*
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡0.359%
Dicamba (3,6-Dichloro-o-Anisic Acid)*** 0.090%
OTHER INGREDIENTS:
TOTAL:
Isomer Specific Method, Equivalent to:
*2,4-Dichlorophenoxyacetic Acid 1.076%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid
***3,6-Dichloro-o-anisic Acid
±CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-₽

EPA REG. NO. 228-401-32802

EPA EST. NO. 59144-GA-1 (A), 32802-MO-1(B), 37351-CO-1 (C) Viper is a Registered Trademark of Nufarm Americas Inc.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, pleae refer to: http://www.epa.gov/espp/litstatus/wtc/index.htm

Distributed for: Howard Johnson's Enterprises, Inc. 9675 S 60th Street, Franklin, WI 53132

Net Weight 50 lbs (22.68kgs) **KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle (If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - PRECAUCION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTION EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

All loaders, applicators, and other handlers must wear:

·long-sleeved shirt and long pants,

•shoes plus socks, and

chemical-resistant gloves

ing/maintaining PPE. If no such instructions for washables exist, or other desirable plants. use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.

• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

FIRST AID

	treatment advice
	 Call a poison control center or doctor for
	first 5 minutes, then continue rinsing eye.
	 Remove contact lenses, if present, after the
	with water for 15 to 20 minutes.
IN EYES	 Hold eye open and rinse slowly and gently

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

This product is a stable mixture of four herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

Not for use on turf grown for sale or other commercial use as sod, or commercial seed production, or for research purposes.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. Discard clothing and other absorbent material that have been DO NOT USE on Dichondra and Carpet grass. DO NOT USE this drenched or heavily contaminated with the product's concentrate. product where desirable clovers are present or allow the product Do not reuse them. Follow manufacturer's instructions for clean- to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees

This product is designed to control many broadleaf weeds such as:

Alder	Crewred is a
Alder	Ground ivy
Annual yellow sweet clover	Gumweed
sweet clover	Hairy bittercress Hairy fleabane
Artichoke	Hairy fleabane
Aster	Hawkweed
Austrian fieldcress	Healall
Bedstraw	Heartleaf drymary
Beggartick	Hedge bindweed
Biden	Hedge bindweed
	Hedge mustard
Bindweed	Hemp
Bird vetch	Henbit
Bitterweed	Hoary cross
Dittor winterere	Hoary cress Hoary plantain
Bitter wintercress	Hoary plantain
Black-eyed Susan	Hoary vervain
Black medic	Honeysuckle
Black mustard	Hop clover
Plackaged plantain	Horsenettle
Blackseed plantain	
Blessed thistle	Horsetail
Blue lettuce	Indiana mallow
Blue vervain	Ironweed
Box elder	Jewelweed
Bracted plantain	Jimsonweed
Brassbuttons	Kochia
Bristly oxtongue	Knawel
Bristly oxtongue Broadleaf dock	Knotweed
Dioauleal uock	
Broadleaf plantain	Lambsquarter
Broomweed	Lespedeza
Buckhorn	Locoweed
Buckhorn plantain	Lupine
	Lupine
Bulbous buttercup	Mallow
Bull nettle	Marshelder
Bull thistle	Matchweed
Burdock	Mexicanweed
Burning nettle	Milk vetch
Bur ragweed	Milkweed bloodflower
Burweed	Mugwort
Buttercup	Morningglory
Canada thiatla	Morninggiory
Canada thistle	Mousear chickweed
Carolina geranium	Musk thistle
Carpetweed	Mustard
Catchweed bedstraw	Narrowleaf plantain
Catagor	Narrowiear plantain
Catsear	Narrowleaf vetch
Catnip	Nettle
Chickweed	Orange hawkweed
Chicory	Oxalis
Cinquefoil	
	Oxeye daisy
Clover	Parsley-piert
Cockle	Parsnip
Cocklebur	Pearlwort
Coffeebeen	Deppyoreae
Coffeebean	Pennycress
Coffeeweed	Pennywort
Common chickweed Common mullein	Peppergrass
Common mullein	Pepperweed
Common sowthistle	Pigwood
	Pigweed Bisouwoodo bodotrow
Corn chamomile	Pineywoods bedstraw
Creeping jenny	Plains coreopsis
Crimson clover	Plantain
Croton	Poison hemlock
Cudwood	Doloon hay
Cudweed	Poison ivy
Curiy dock	Poison oak
Curly dock Curly indigo	Pokeweed
Dandelion	Poorjoe
Dead nettle	Povertyweed
	Prickly lottuco
Dock	Prickly lettuce Prickly sida
Dogbane	Prickly sida
Dogfennel	Primrose
Elderberry	Prostrate knotweed
English daisy	
Fall deadalian	Prostrate pigweed
Fall dandelion	Prostrate spurge
False dandelion	Prostrate vervain
False flax	Puncture vine
False sunflower	Purslane
Fiddlepook	Dogwood
Fiddleneck	Ragweed Red clover
Field bindweed	Red clover
Field pansy	Redroot pigweed
Field pansy Flea bane (daisy)	
Eliverood	Red sorrel
Flixweed	Redstem filaree
Florida betony	Rough cinquefoil
Florida pusley	Rough fleabane
Frenchweed	Roundleafed marigold
Calineana	
Galinsoga	Rush
Garlic mustard	Russian pigweed
Goathead	Russian thistle
Goatsbeard	St. Johnswort
e catobourd	

Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsog Smartweed Smooth dock Smooth pigweed Sneezev Southern wild rose Sowthistle Spanishneedle Spatterdock Spiny Amaranth Spiny cocklebur Spotted catsear Spotted knapwee Spotted spurge Spurge Spurweed Stinging nettle Stinkweed Stitchwort Strawberry clover Sumac Sunflower Sweet clover Tall nettle Tall vervain Tansy mustare Tansy ragwor Tanwee Tarweed Thistle Tick trefoil Toadflax Trailing Crownvetch Tumble mustard Tumble pigweed Tumblew Velvet leaf Venice mallow Veronica Vervain Vetch Virginia buttonweed Virginia creeper Virginia pepperweed Wavyleaf bullthistle Western clematis Western salsify White clover White mustard Wild mustard Wild aster Wild buckwheat Wild carrot Wild four-o'-clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potat Wild vetch Willow Witchweed Wooly morningglor Woodsorrel Woolv croton Wooly plantair Wormseed Yarrow Yellow rocket Yellowflower pepperweed, and other broadleaf weeds

ing actively in Spring or early Fall. Avoid contact with desirable snrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLshrubs, plants, vegetables, or flowers; especially when applying with weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT retreatment in 30 days may be necessary if lawn is heavily overrun Liable to BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSwith weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, overapplication may cause minor transitory discoloration on Centipede If you do not agree with or do not accept any of directions for use, the warranty disand St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 15,625 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide, Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.8 pounds per 250 square feet.

Spreader settings to apply at listed rate 3.2 lbs. per 1000 sq.			
Scotts Accu-Pro 2000	М		
Scotts SR2000	М		
Spyker 76-22	4		
Lesco 021820	H(18)		
PSB Bigfoot	J		

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion

Limitations: The maximum rate per application is 3.2 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 6.4 pounds per 1,000 square feet, excluding spot treatments. Minimum of 30 days between applications.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed, or other pesticides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office or quidanc

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse of refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sani tary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CON-SISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTH-ERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF WHEN TO APPLY: Best results are obtained when weeds are grow-ing actively in Spring or early Fall Avoid contact with desirable INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINA ER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRE-SENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY

TO THESE GOODS. LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFAC TURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAM-AGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BYTHE BUYER, USER, OR ITS CUSTOMERS TOTHE EXTENT CON-SISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MAN UFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEED ING THE COST OF THE PRODUCT

claimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. This product is protected by U.S. Patent No. 6,579,831. No license granted under any nor U.S. Patent(s)