

HJ

Howard Johnson's
WEED & FEED
 CONTAINS
SURGE® BROADLEAF HERBICIDE

Not for sale, distribution or use in Nassau or Suffolk Counties in New York State

ACTIVE INGREDIENTS:
 2,4-D, 2-ethylhexyl ester1.153%
 Mecoprop-p acid0.174%
 Dicamba acid0.073%
 Sulfentrazone0.034%
 OTHER INGREDIENTS:98.566%
 TOTAL100.000%

THIS PRODUCT CONTAINS:
 15.30 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.765% (CAS 1928-43-4)
 3.48 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.174% (CAS 16484-77-8)
 1.46 lbs. 3,6-dichloro-o-anisic acid equivalent per ton or 0.073% (CAS 1918-00-9)
 0.68 lbs. N-[2,4-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]phenyl] methanesulfonamide per ton or 0.034% (CAS 122836-35-5)
 Isomer Specific by AOAC Methods
 Surge® are registered trademarks of PBI/Gordon Corp.
 EPA Reg. No. 2217-882-32802
 EPA EST. NO. 59144-GA-1 (A), 32802-MO-1(B), 37351-CO-1 (C)

Distributed by: **Howard Johnson's Enterprises, Inc.**
 9675 S 60th Street, Franklin, WI 53132

KEEP OUT OF REACH OF CHILDREN

CAUTION

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)
 All loaders, applicators, and other handlers must wear:
 • long-sleeved shirt and long pants,
 • shoes plus socks,
 • waterproof gloves

User Safety Requirements
 Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

User Safety Recommendations
 • Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
 • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 • If pesticide gets on skin, wash immediately with soap and water.
 • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards
 This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate. This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

First Aid	
If in eyes:	• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment information.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

16-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N)16%
 16.00% Urea Nitrogen*
 Soluble Potash (K₂O)4%
 Sulfur (S)1.36%
 1.36% Combined Sulfur (S)

Derived From: Polymer Coated Urea, Sulfate of Potash, Urea

*3.20% Slowly Available Nitrogen from Polymer Coated Urea

Chlorine (Cl), not more than6.00%

F315
 100526094

Net Weight 50 Lbs (22.68 kg)

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
USE RESTRICTIONS:
 • Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
 • Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
 • Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers.
 • Do not apply this product to newly seeded grasses until well-established or after the third mowing.
 • Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area.
 • Do not use this product on 'Floritam' St. Augustinegrass.
 • Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.

STATE RESTRICTIONS
 Keep this product off sidewalks, patios, driveways, etc. This fertilizer may contain iron which will stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.
 Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street, back into your lawn/garden.

WHERE TO USE:
 Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide is a herbicide containing fertilizer, 2,4-D, mecoprop-p, dicamba and sulfentrazone. This granular weed and feed provides plant nutrients and controls a broad spectrum of listed broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.
 Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide is intended to be applied by noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example: hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide may be applied to newly seeded cool season turfgrass. Apply only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the listed rate for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area. Do not apply product to bare ground. Do not use this product on dichondra, carpetgrass, creeping bentgrass, or turf containing desirable clovers.

California: Make broadcast applications only between March 1 and September 1. While irrigation is necessary and important for plant growth, apply irrigation water efficiently so that no more than 125% of the net irrigation requirement is applied for any irrigation event. Apply efficient irrigations for six months following application of sulfentrazone containing products.

New York: Only one application per year of this product is allowed. This product is not allowed to be sold or used in Nassau or Suffolk Counties.

Limitations on broadcast applications to ornamental turfgrass.				
Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Seasonal Rate
Ornamental turfgrass	174 pounds/A (1.33 lb 2,4-D ae/A) (0.30 lb MCPP-p ae/A) (0.13 lb dicamba ae/A)	2	30 Days	348 pounds/A (2.66 lb 2,4-D ae/A) (0.60 lb MCPP-p ae/A) (0.25 lb dicamba ae/A)

ae= Acid Equivalent. Do not exceed the maximum seasonal rate of 348 pounds (2.66 lb 2,4-D ae), 0.60 lb MCPP-p ae and 0.25 lb dicamba ae) per acre per season, excluding spot treatments.

HOW MUCH TO APPLY:
For Kentucky bluegrass, fescue spp., perennial ryegrass, common bermudagrass, and zoysiagrass:
 Apply 130 to 174 pounds of Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide per acre of established turfgrass. The equivalent application rates are 3.0 to 4.0 pounds of product per 1,000 square feet of turfgrass.

For New York State only:
 Apply 130 to 166 pounds of Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide per acre of established turfgrass. The equivalent application rates are 3.0 to 3.8 pounds of product per 1,000 square feet of turfgrass.

For St. Augustinegrass, centipedegrass, hybrid bermudagrass, bentgrass, and colonial bentgrass:
 Apply 87 to 109 pounds of Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide per acre of established turfgrass. The equivalent application rates are 2.0 to 2.5 pounds of product per 1,000 square feet of turfgrass.

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

APPLICATION DIRECTIONS:
 1. Spring and fall applications are suitable for the best control. However, Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide can be applied anytime broadleaf weeds are present and actively growing.
 2. Mow the turfgrass 1 to 2 days before application.
 3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
 4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
 5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

Bag Size,Lbs.	Application Rate,Lb/1,000 Sq.Ft.	Treated Area for This Bag Size, Sq.Ft.	
50	2.0	25,000	
	2.5	20,000	
	3.0	16,700	
	3.2	15,600	
	3.4	14,700	
	3.6	13,900	
	4.0	12,500	

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer. Refer to the rate table for the appropriate bag sizes and the area to be treated.

SPREADER	Delivers 4 lb /1,000 sq.ft.	
	Cone Setting	4 lbs
Accu-Pro	6	M 1/2
SR2000	6	N
Lesco	0.25 closed	E 1/2
Spyker 76-22	4.5	
PSB Bigfoot	A.5	K 1/2
Lely HR	II-B	6
Vicon		17

CALIBRATION INSTRUCTIONS:
 To provide proper distribution, calibrate spreader before application.
 For 4.0 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound of product per 250 square feet.

DISTRIBUTION AND SPREADER PATTERNS:
 Always apply with a properly calibrated rotary or drop type fertilizer spreader to distribute the granules evenly. Never apply by hand.

Coverage - To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One 50 pound bag covers up to 12,500 sq. ft.

SINGLE PASS DISTRIBUTION:
 For best results, apply this product evenly and uniformly avoiding streaking, skips or overlaps.
 1. For rotary spreaders, operate at intervals corresponding to the swath width. For drop spreaders, overlap the wheel tracks to ensure complete coverage.
 2. First, spread the borders of the lawn. Then apply this product to the center with straight line passes.
 3. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

WEEDS CONTROLLED:
 Dandelion, chickweed, plantain, and clover are controlled effectively with Howard Johnson's Weed & Feed Contains Surge Broadleaf Herbicide. A list of susceptible weeds is shown below.

- BROADLEAF WEEDS -
 Annual yellow sweetclover, Aster, Austrian fieldcress, Bedstraw, Beggarticks, Betony, Florida, Bird vetch, Bitter wintercress, Bittercress; hairy, Bitterweed, Black-eyed Susan, Black medic, Blackseed plantain, Blessed thistle, Bloodflower milkweed, Blue lettuce, Blue vervain, Bracted plantain, Bristly oxtongue, Broadleaf dock, Broadleaf plantain, Buckhorn plantain, Bulbous buttercup, Bull thistle, Bullnettle, Burclover, Burdock, Burning nettle, Burweed, Buttercup, Buttonweed, Canada thistle, Carolina geranium, Carpetweed, Catchweed bedstraw, Catnip, Catsear, Chickweed; common, Chickweed; mouseear, Chicory, Cinquefoil, Clover; crimson, Clover; hop, Clover; red, Clover; strawberry, Clover; sweet, Clover; white, Cockle, Creeping jenny, Cudweed, Curly dock, Daisy; English, Daisy fleabane, Daisy; oxeye, Dandelion, Dichondra, Dollarweed, False dandelion, Falseflax, False sunflower, Florida pusley, Galinsoga, Goldenrod, Ground ivy, Gumweed Hairy fleabane, Hawkweed, Healall, Heartleaf drymary, Heath aster, Henbit, Hoary cress, Hoary plantain, Hoary vervain, Knawel, Knotweed, Lambsquarters, Lespedeza, Mallow, Matchweed, Mexicanweed, Milk vetch, Mouseear hawkweed, Mugwort, Narrowleaf plantain, Nettle, Orange hawkweed, Oxalis Parsley-piert, Parsnip, Pearlwort, Pepperweed, Pigweed, Pinewoods bedstraw, Plains coreopsis, Plantain, Poison ivy, Poison oak, Poojoe, Prairie sunflower, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Puncturevine, Purslane; common, Ragweed, Red sorrel, Redroot pigweed, Redstem filaree, Rough cinquefoil, Russian pigweed, Scarlet pigweed, Scarlet pimpernel, Sheep sorrel, Shepherdspurse, Slender plantain, Smallflower galinsoga, Smooth dock, Smooth pigweed, Sorrel, Sowthistle, Spanishneedles, Speedwell, Spiny amaranth, Spiny sowthistle, Spotted catsear, Spotted spurge, Spurweed, Strawberry; India mock, Tall nettle, Tansy ragwort, Tansy mustard, Thistle, Trailing crownvetch, Tumble mustard, Tumble pigweed, Venice mallow, Virginia buttonweed, Virginia pepperweed, Wavyleaf bullthistle, Western clematis, Wild aster, Wild carrot, Wild four-o'clock, Wild garlic, Wild geranium, Wild lettuce, Wild marigold, Wild onion, Wild rape, Wild strawberry, Wild vetch, Woodsorrel, Woolly croton, Woolly plantain, Wormseed, Yarrow, Yellow rocket, Yellowflower pepperweed

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.
PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.
PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.
CONTAINER HANDLING: Nonrefillable container: Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

LIMITED WARRANTY AND DISCLAIMER
 FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.