

Viper® Granular **Weed Control**

ACTIVE INGREDIENTS:
Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid* 1.346%
2,4-Dichlorophenoxyacetic Acid*
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡
Dicamba (3,6-Dichloro-o-Anisic Acid)***
OTHER INGREDIENTS:
TOTAL :
Isomer Specific Method, Equivalent to:
*2,4-Dichlorophenoxyacetic Acid
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid
***3,6-Dichloro-o-anisic Acid
‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p
TCONTAINS THE SINGLE ISOMER FORM OF MECOPROP-P

EPA REG. NO. 228-401-32802

EPA EST. NO. 59144-GA-1 (A), 32802-MO-1(B), 37351-CO-1 (C)

Viper is a Registered Trademark of Nufarm Americas Inc.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in <u>Washington Toxics</u> water or groundwater. <u>Coalition, et al. v. EPA</u>. C01-0132C, (W.D. WA). For further information, please refert to: http://www.epa.gov/espp/litstatus/wtc/index.htm

Distributed for: Howard Johnson's Enterprises, Inc. 9675 S 60th Street, Franklin, WI 53132

Net Weight 20 lbs (9.07 kgs) **KEEP OUT OF REACH OF CHILDREN CAUTION - PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a

(If you do not understand the label, find someone to explain it to you in detail.)

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

PRECAUTIONARY STATEMENTS **CAUTION - PRECAUCION**

thoroughly wash hands and exposed skin with soap and water.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF IN EYES

- ·Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- •Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS

This product is a stable mixture of four herbicides. It is for use on residental turf.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish HAZARDS TO HUMANS AND DOMESTIC ANIMALS ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to Causes moderate eye irritation. Avoid contact with skin, eyes, or your lawn or garden, and sweep any product landing on the driveclothing. Wear Long pants, long-sleeved shirt, socks and shoes way, sidewalk, gutter, or street back onto the treated area. To prewhen applying this product. After using this product, remove vent product run-off, do not over water the treated area(s) to point clothing and launder separately before reuse, and promptly and of runoff or apply when raining or when rain is expected that day.

The maximum number of applications per treatment site is 2 per

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass.

DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to control many broadleaf weeds such as:

Scotch thistle

Annual yellow sweet clover Artichoke Hairy fleabane Slender plantain Aster Hawkweed Smallflower galinsog Austrian fieldcress Bedstraw Beggartick Biden Healall Heartleaf drymary Smartweed Smooth dock Hedge bindweed Hedge mustard Smooth pigweed Sneezeweed Bindweed Southern wild rose Sowthistle Bird vetch Bitterweed Hoary cress Spanishneedle Bitter wintercress Hoary plantain Hoary vervain Spatterdock Black-eyed Susar Black medic Black mustard Spiny Amaranth Honeysuckle Hop clover Spiny cocklebur Blackseed plantair Spotted catsear Horsetail Blessed thistle Spotted knapwee Blue lettuce Indiana mallov Spotted spurge Blue vervain Ironweed Spurge Spurweed Box elde Bracted plantain Brassbuttons Kochia Knawel Stitchwort Bristly oxtonau Broadleaf dock Knotweed Strawberry cloves Broadleaf plantai Lambsquarter Lespedeza Buckhorn Locoweed Sweet clover Buckhorn plantair Tall nettle Marshelder Bull thistle Matchweed Tansy ragwor Burdock Mexicanweed Burning nettle Milkweed bloodflowe Bur ragweed Thistle Tick trefoil Burweed Muawort Morningglory Mousear chickweed Musk thistle Toadflax
Trailing Crownvetch
Tumble mustard Carolina geraniun Mustard Narrowleaf plantain Narrowleaf vetch Carpetweed Tumble pigweed atchweed bedstra Tumblew Catnip Venice mallow Chickweed Orange hawkweed Veronica Oxalis Oxeye daisy Vetch Virginia buttonweed Clover Parsnip Pearlwort Pennycress Virginia creeper Virginia pepperweed Wavyleaf bullthistle Coffeebean Coffeeweed Pennywort Western clematis Common chickweed Common mullein Common sowthistle Corn chamomile Peppergrass
Pepperweed
Pigweed
Pineywoods bedstraw Western salsify Wild mustard Plains coreopsis Creeping jenny Crimson clover Wild aster Plantain Wild buckwheat roton Poison hemlock Wild carrot Wild four-o'-clock Cudweed Poison ivy Poison oak Curly dock Curly indigo Dandelion Dead nettle Wild garlic Pokeweed Poorjoe Povertyweed Wild marigold Dock Prickly lettuce Wild onion Dock Dogbane Dogfennel Elderberry English daisy Prickly sida Wild parsnip Wild radish Prostrate knotweed Wild rape Prostrate pigweed Prostrate spurge Prostrate vervain Wild strawberry Wild sweet potato Wild vetch Fall dandelio False dandelion False flax Puncture vine Willow Witchweed False sunflowe Purslane Fiddleneck Field bindweed Field pansy Flea bane (daisy) Redroot pigweed Wooly croton Red sorrel Redstem filaree Rough cinquefoil Wooly cloton Wooly plantain Wormseed Yarrow Flixweed Florida betony Yellow rocket Florida pusle Rough fleabane Frenchweed Roundleafed marigol Galinsoga Garlic mustard Goathead Rush Russian pigweed Russian thistle Goatsbeard St .lohnswort

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable ER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPREshrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible TOMEDS OF BLYER IS ANY OF INDEMNIFICATION TO BUYER OR TO CUSretreatment in 30 days may be necessary if lawn is heavily overrun DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF with weeds or if adverse low moisture conditions prevail and weeds THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, are in a state of poor growth. Do not exceed specified rates because OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS damage to turf may occur.

inadequate moisture, high temperature and high humidity, over- ING THE COST OF THE PRODUCT. application may cause minor transitory discoloration on Centipede If you do not agree with or do not accept any of directions for use, the warranty disand St. Augustine grasses. It is recommended that during hot, dry seller, and the purchase price will be refunded. conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 6,250 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2 pounds per 1,000 square feet. These settings are furnished only as a guide, Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.8 pounds per 250 square feet.

Spreader settings to apply at listed rate 3.2 lbs. per 1000 sq.	
Scotts Accu-Pro 2000	M
Scotts SR2000	M
Spyker 76-22	4
Lesco 021820	H(18)
PSB Bigfoot	J

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion

Limitations: The maximum rate per application is 3.2 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 6.4 pounds per 1,000 square feet, excluding spot treatments. Minimum of 30 days between applications.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. PESTICIDE STORAGE: Store in a cool, dry place. Keep container closed and away from food, feedstuffs, and domestic

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CON-SISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTH-ERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINA-TION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELL-SENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. IN NO EVENT SHALL MANUFAC-TURER OR SELLER BE LIABLE FOR SPECIAL INCIDENTAL OR CONSEQUENTIAL DAM-AGES. OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW. MANUFACTURER OR SELLER SHALL NOT BE TOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CON-SISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MAN-NOTE: When applying to Southern Grasses - under conditions of UFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEED-

claimers, or limitations on liability, do not use the product, and return it unopened to the

(RV011615) 228-401 Nufarm, retail 01-16-15 (hie100818)