



All-Season

WEED & FEED

With Lazer®

Controls a wide spectrum of broadleaf weeds.



18·0·3

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-D Acid*	0.664%
Dimethylamine Salt of Mecoprop-p Acid**‡	0.163%
Dicamba Acid***	0.028%
OTHER INGREDIENTS:	99.145%
TOTAL:	100.000%

Isomer Specific Method, Equivalent to:

* 2,4-D Acid	0.552%
** Mecoprop-p Acid	0.135%
*** Dicamba Acid	0.028%


‡ CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-P

EPA Reg. No. 228-432-32802
EPA Est. No. 59144-GA-1 (A)
EPA Est. No. 32802-MO-1 (B)
EPA Est. No. 37351-CO-1 (C)

Lazer is a Registered Trademark of Nufarm Americas Inc.

Covers 5,000 sq. ft. Net Weight: 16 lbs. (7.26 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION: SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS



WARRANTY DISCLAIMER
The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE LIMITATION OF LIABILITY, AND (2) MANUFACTURER OR SELLER SHALL BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDemnIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of the directions for use, the warranty disclaimer, or limitations on liability, do not use the product, and return it unopened to the seller, and the purchase price will be refunded. (R202710)

© Copyright 2011 Howard Johnson's Enterprises, Inc.
9675 S 60th Street, Franklin, WI 53132

Do not contaminate water, food, or feed by storage or disposal.
PESTICIDE STORAGE: Always use original container to store pesticides and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides.
Do not stack more than two pallets high.
CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

Drop Spreader	Broadcast Spreader	Agri-Fab	Ames	Ames Deluxe 2in1	Craftsman	Earthway	Plant Mate	Precision GreenLine	Precision Red Devil	Precision Pro	Scotts Speedy Green	Scotts Easy Green	Scotts Basic	Scotts Lawn Pro	Scotts Deluxe EdgeGuard	Taylor
4.0	4.0	4.0	3.0	1.5	4.0	1.5	8.0	3.75	3.75	3.75	5.75	26.0	5	5	5	8.0
6.0	4.0	4.0	4.0	1.5	5.5	9.5	1.7	1.5	1.5	1.5	6.5	7.5	1.7			

Drop Spreader
Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

The following settings are approximate for applying contents of this package at a suggested rate of 3.2-2.2 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 0.8 pound per 250 square feet.

How to Apply

Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinning) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results: (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment.

When to Apply

Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

WEEDS CONTROLLED

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Alder, Annual yellow sweet clover, Artichoke, Aster, Australian fieldcress, Bedstraw, Beggartick, Bidens, Bindweed, Bird vetch, Bitterweed, Bitter wintercress, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Blue lettuce, Blue vervain, Box elder, Bracted plantain, Brassbutton, Britly ox-tongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common chickweed, Common mullen, Common carpweed, Carpetweed, Catchweed bedstraw, Catsear, Catnip, Chickweed, Chicory, Cinqufoil, Clover, Burdock, Burning nettle, Bur ragweed, Burweed, Buttercup, Canada thistle, Carolina geranium, Plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Blue vervain, Box elder, Bracted plantain, Brassbutton, Britly ox-tongue, Broadleaf dock, Broadleaf plantain, Broomweed, Buckhorn, Buckhorn plantain, Bulbous buttercup, Bull nettle, Bull thistle, Black-eyed Susan, Black medic, Black mustard, Blackseed plantain, Blessed thistle, Blue lettuce, Stinkweed, Stitchwort, Strawberry clover, Sumac, Sunflower, Sweet clover, Tall nettle, Tall vervain, Tansy mustard, Tansy ragwort, Tanweed, Thistle, Tick trefoil, Toadflax, Trailing Crownvetch, Cocklebur, Spotted catsear, Spotted knapweed, Spotted spurge, Spurge, Spunweed, Stinging nettle, Southern wild rose, Sowthistle, Spanshneedle, Spatterdock, Speedwell, Spiny Amaranth, Spiny plantain, Smallflower gallinules, Smartweed, Smooth dock, Smooth pigweed, Sneezeweed, thisle, St. Johnswort, Scarlet pimpernel, Scotch thistle, Sheep sorrel, Shepherdspurse, Slender vetch, Rough cinquefoil, Rough fescue, Roundleafed margold, Russian pigweed, Russian Puncture vine, Purslane, Ragweed, Red clover, Redroot pigweed, Red sorrel, Redstem flaxweed, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poortoe, Povertyweed, Prickly lettuce, Prickly sida, Primrose, Prostrate knotweed, Prostrate pigweed, Prostrate spurge, Prostrate vervain, Parnwort, Peppergrass, Peppercorn, Pigweed, Pinewoods bedstraw, Plains coreopsis, Plantain, vetch, Nettle, Orange hawkweed, Oxalis, Oxye daisy, Parsley-piert, Parsnip, Pearlwort, Pennygrass, Mugwort, Morningglory, Mouseear chickweed, Musk thistle, Mustard, Narrowleaf plantain, Narrowleaf Lupine, Mallow, Marsbaler, Matchweed, Mexicanweed, Milk vetch, Milkweed, Bloodflower, Jeweled, Jimsonweed, Kochia, Knawel, Knotweed, Lambsquarter, Lespedeza, Locoweed, plantain, Hoary vervain, Honeybuckle, Hop clover, Horehound, Horsetail, Indiana mallow, Hoary Heall, Heartleaf drymary, Hedge bindweed, Hedge mustard, Hemp, Henbit, Hoary cress, Hoary Goatsbeard, Goldenrod, Ground ivy, Gunweed, Hairy bittercress, Hairy fleabane, Hawkweed, (galxy), Flaxweed, Florida pusley, Florida pusley, Frenchweed, Gallinsoa, Garlic mustard, Goathead,

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water; also before eating, drinking, chewing gum or using tobacco.
ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.
 This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product is a stable mixture of three herbicides. It is for use on residential turf.
 This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grasses. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

Do not apply more than 2 applications per year, including broadcast applications in combination with spot treatments.
 Do not enter treatment area until dust has settled.
 Do not apply to treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment.

GENERAL PRECAUTIONS AND RESTRICTIONS:

The maximum number of applications per treatment site is 2 per year. Do not allow people (other than the applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

Directions for Use

For Medical Emergencies Only, Call (877) 325-1840. For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300.

treatment information.
HOT LINE NUMBER Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

IF SWALLOWED
 • Call a poison control center or doctor immediately for treatment advice.
 • Have person sip a glass of water if able to swallow.
 • Do not induce vomiting unless told by a poison control center or doctor.
 • Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING
 • Take off contaminated clothing.
 • Rinse skin immediately with plenty of water for 15-20 minutes.
 • Call a poison control center or doctor for treatment advice.

IF IN EYES
 • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
 • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
 • Call a poison control center or doctor for treatment advice.

FIRST AID

Precautionary Statements

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfc.org/metals.htm>

Total Nitrogen (N)..... 18.00%
 18.00% Urea Nitrogen
 Soluble Potash (K₂O)..... 3.00%
 Derived From: Muriate of Potash, Urea
 Chlorine (Cl), not more than 6.00%
 Fertilizer guaranteed by Howard Johnson's Enterprises, Inc.
 F315

Guaranteed Analysis

For a lush green lawn
 Vital for growth & drought resistance

3
POTASH

18
NITROGEN